

Federal Food, Drug and Cosmetic Act

A food is adulterated:

- If it bears or contains any poisonous or deleterious substance which may render it injurious to health, or
- If it bears or contains any added poisonous or deleterious substance.

Chemical Hazards

Ways in which the hazard is introduced into the facility.

- Incoming Raw Ingredients
- Employee Practices
 - ▶ Batching
 - ➤ Mixing
 - > Warehousing
 - ▶ Load out

Mytocoxins developed in the field or storage



Laws and Regulations

- □ Require that foods are safe and free of adulterants.
- The FDA, EPA and USDA enforce provisions of pertinent laws and regulations

Pesticide Contamination



1

Cross Contamination during Storage or Processing



Fungal Toxins

Fungus	Toxin
Aspergillus flavus &	aflatoxin
A. parasiticus	
Fusarium species	tricothecenes
F. Verticillioides	fumonisins
F. Graminearum	zearalenone
Penicillium verrucosum	ochratoxins
Claviceps	ergot

Contamination from Equipment Carryover



Drug Residues

- Incoming ingredients
 - DDGS
- Cross Contamination
 - Mixer
 - Conveyors
 - Holding bins
- Weighing out wrong ingredient
 - Hand adds
 - Charging micro-bin with wrong ingredient
 - Formulation error

Types of Chemical Hazards

Mycotoxins

Produced in feeds as byproducts from the growth of specific types of molds

Drug residues

- Unapproved
- Exceeding Tolerance levels

Toxic Elements/Compounds

Dioxin and PCBs

Pesticide Residues

Heavy Metals

- Lead (Pb) Lead is a naturally-occurring neurotoxin. Lead toxicity primarily targets the nervous system, kidneys, bones, heart and blood, and poses greatest risk to infants, young children and pregnant women.
- Mercury (Hg) Mercury toxicity can affect the central nervous system, kidneys and liver.
- Arsenic (As) Arsenic toxicity seems to predominantly affect the skin, lungs and gastrointestinal system, and may cause nervous disorders, deteriorated motor coordination, respiratory diseases, and kidney damage as well as cancers of the skin, liver, bladder and lungs.
- Cadmium (Cd) primarily affect the lungs, kidneys, bones, and immune system
- Others:

http://otscweb.tamu.edu/OTSC-Present/2011/Heavy%20Metal%20Analysis-10-21-2011-wei.pdf

THE TEXAS STATE CHEMIST

Pesticide Residues

 Dyfonate Case Study: Found in meat and bone meal in Texas at 0.86 ppm

EPA tolerances for this organophosphate insecticide that were located in 40 CFR 180.221. These tolerances were established for O-Ethyl S-phenyl ethylphosphonodithioate, which is the chemical name for fonofos, and ranged from 0.5 ppm on asparagus to 0.1 ppm on several raw agricultural commodities including corn grain and peanuts. All these tolerances were revoked as of December 31, 2002 so today any quantifiable level of fonofos on a feed ingredient is a technical violation.

Control of Chemical Hazards

Products received into the facility

- Approved Suppliers
- Pre-Requisite Programs
 - ➤ Visual Inspection
 - > Sampling and analysis

During Processing

- Employee Practices
- Prerequisite Programs

OFFICE OF THE TEYAS STATE CHEMIST

Dioxin and PCBs

- Sources: clay, citrus pulp, fish meal and oil, other sources of fat and oil, trace metal ingredients manufactured through the smeltering process
- Testing: High resolution GC, high resolution mass spectrometry, bioassay technology for screening
- Recent Incidents: German Agriculture Minister Ilse Aigner on Tuesday urged state officials to implement a federal action plan that aims to prevent future food contamination ahead of special government summit on the dioxin animal feed scandal. (Jan 21, 2011).
 - The EU pig market is in crisis due to high cereal prices and the dioxin problems in Germany.
- Hazard: wasting, skin disorders, immune suppression, enlarged liver, endocrine disruption

Summary

Chemical hazards exist in every facility.

HACCP Teams must ask themselves

- What is the likelihood of the hazard?
- Do we already have something in place to control that hazard?

Types of Chemical Hazards

Allergens (tend not to be a concern in livestock feeds)

- Proteins that induce an allergic reaction.
- Symptoms range from mild rashes to anaphylaxis and, rarely, death.
- 1-2% of adults have food allergy.
- 6% of children have a food allergy.

END

Dr. Tim Herrman
Professor, State Chemist & Director
Texas A&M University
(979) 845-1121
tjh@otsc.tamu.edu

OFFICE OF THE TEXAS STATE CHEMIST
Texas Feed and Fertilizer Control Service & Agriculture Analytical Service

3